

Article

Impact of Seed Transmission on the Infection and Development of Pitch Canker Disease in *Pinus radiata*

Margarita Evira-Recuenco ¹, Eugenia Iturritxa ² and Rosa Raposo ^{1,*}

¹ Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria, Centro de Investigación Forestal (INIA-CIFOR), Madrid 28040, Spain; E-Mail: elvira@inia.es

² NEIKER, Granja Modelo de Arkaute. Apdo 46. Vitoria-Gasteiz 01080, Spain; E-Mail: eiturritxa@neiker.eus

* Author to whom correspondence should be addressed; E-Mail: raposo@inia.es; Tel.: +34-91-347-3954; Fax: +34-91-347-6883.

Academic Editors: Jan Stenlid, Jonas Oliva and Audrius Menkis

Received: 25 August 2015 / Accepted: 17 September 2015 / Published: 23 September 2015

Abstract: The aim of this study was to determine the impact of seed inoculum on subsequent disease development in nurseries, specifically studying incidence of seed infection, transmission rate of seed to seedlings, and rate of disease development from the primary inoculum source. Transmission rate of *Fusarium circinatum* (*F. circinatum*) from symptomatic trees of *Pinus radiata* (*P. radiata*) to seed was 0.73%, being the fungus mainly on the coat. Seed infection incidence was positively correlated with tree disease severity. Seeds also become contaminated with *F. circinatum* during storage, where high relative humidity had a great effect regardless of temperature. Transmission rates from seeds to symptomatic seedlings measured at 48 days after seed inoculation with 10^4 and 10^6 was 28% and 80%, respectively. Seedlings showed symptoms of wilting and dieback, and eventually died within the total duration of this study (475 days). By this time, there were two remaining symptomless seedlings (1% of the total). *F. circinatum* was recovered from each of them, demonstrating its persistence in seedlings. Fungus populations in roots were estimated between 10^5 and 10^7 cfu/g in symptomatic plants for both inoculum doses and up to 10^4 cfu/g in asymptomatic plants. Disease infection rate was similar in post-emergence damping-off (up two months growing) and for established seedlings (up to eight months) after seeds were inoculated with 10^4 spores/mL.

Keywords: *Fusarium circinatum*; nursery; seed infection incidence; seed contamination; seed transmission rate; persistence; infection rate; damping-off

1. Introduction

Pitch canker disease, caused by *Fusarium circinatum* (*F. circinatum*) Nirenberg and O'Donnell, is an emergent disease that is threatening *Pinus* trees in Europe. Since it was first recognized in Southeastern USA [1], the disease has been reported in other parts of the world [2]. In many of these reports, the disease appeared first in forest nurseries, like in South Africa [3], Chile [4], Spain [5], Portugal [6] and Brazil [7]. In trees, *F. circinatum* causes branch dieback and resinous cankers that progressively girdle the wood causing branch death. Canker infection of multiple branches of the main stem can cause extensive dieback in the tree crown and tree mortality.

Pine seeds are considered one of the major pathways of introduction of *F. circinatum* into new countries, as it occurred in Chile [8], South Africa [9], and Spain [10]. Pine seed is also one of the potential inoculum sources of *F. circinatum* in nurseries [2,11,12]. This fungus is a seed-borne pathogen in *Pinus radiata* [11]; it has also been detected in seeds of *P. taeda* [13] and *P. elliottii* [14]. *Fusarium circinatum* may infest seeds superficially in cones growing on healthy branches [11]. The fungus can also infect seeds internally, where it causes damage to embryo and gametophyte tissues [13]. The mechanisms by which *F. circinatum* infects seed are unknown, but three types of infection have been identified [11]: superficial, active internal and dormant internal, though the latter is not detectable on plate medium. The extent of internal and/or external seed contamination depends on the pine species and environmental conditions [2]. *Fusarium circinatum* was isolated from up to 83% of seeds of *P. radiata* [11]. Anderson *et al.* [15] determined that 34% of *P. elliottii* infested seed was internal.

In forest nurseries, *F. circinatum* causes pre- and post-emergence damping-off of seedlings [11] and mortality of established seedlings [3]. Main symptoms are collar rot and wilting [2], but under certain conditions, seedlings can be asymptomatic and may develop the disease after they are transplanted to plantations [2,11]. From an epidemiological point of view, host growth stage is an important parameter influencing disease intensity that should be taken into account for disease management [16]. Viljoen *et al.* [3] found that mortality of established seedlings tends to be lower than that of newly germinated seedlings, and Storer *et al.* [11] determined that seedling emergence from infested Monterey pine seed was 9% compared with 67% for non-infested seeds. Another important epidemiological parameter is the infection rate [16], for which there are no available data, comparing infection of newly germinated seedlings and established seedlings.

Little is known about this pitch canker disease in forest nurseries in Spain. *Fusarium circinatum* may be detected in any of the usual species found in these nurseries (*P. pinaster*, *P. sylvestris*, *P. nigra* and *P. radiata*) [5,17], although disease has only been recorded in plantations of *P. radiata* [18,19] and recently, in *P. pinaster* [19]. Disease outbreaks in nurseries are sporadic, which suggests an external source of inoculum. On the other hand, routine seed health tests performed following the specific method for the detection of *F. circinatum* on *Pinus* spp. as recommended by ISTA [20] reveal that number of seeds with presence of the pathogen is almost null. This standard procedure is based on placing seeds on

semi-selective media, and it does not distinguish internal from external presence of *F. circinatum*. Moreover, the presence of the fungus does not indicate necessarily that it has been transmitted from tree to seed, since fungus could have contaminated seeds during commercial processes of harvesting, cleaning and storage. The objectives of this work are to (1) determine the frequency of seeds with presence of *F. circinatum* collected from *P. radiata* trees, and evaluate the importance of seed contamination during its storage; (2) study persistence of the pathogen in seedlings, to demonstrate transmission of seed borne inoculum to the seedling; (3) study the effect of seed inoculum location and dose on infection and development of different host growth stages, including germination; and (4) perform a temporal analysis of the disease with different doses of inoculum. These evaluations will be performed under optimal conditions of temperature [21] on *P. radiata*, one of the most susceptible species [18,22], to determine the maximum disease infection rate.

2. Experimental Section

2.1. Natural Presence of *Fusarium circinatum* in/on Seeds of Symptomatic Trees

In eight plantations of *P. radiata* located in Northern Spain (País Vasco), cones on damaged stems and branches were collected in 2009 from each of 10 trees with symptoms of pitch canker disease. Disease was visually evaluated by the number of cankers and percentage of defoliation. Cones from each tree were pooled and then 100 seeds were obtained by exposing cones to a temperature of 35 °C–40 °C for 24 h to cause them to open. Seeds (after hand-removing their wings) were dissected into separate fractions consisting of seed coat, gametophyte and embryo, following the protocol described by Aitken and Iturrutxa [23] and then cultivated separately in FSM (Fusarium Selective Medium) [24]. Presence of *Fusarium circinatum* was evaluated after incubation for 14 day at 20 °C ± 2 °C.

Logistic regression was used to analyze the relation between the proportion of seeds with the pathogen and the number of cankers per tree (as a measure of severity disease). All statistical analyses were conducted using SAS, Version 9.3 (SAS Institute Inc., Cary, NC, USA).

2.2. Seed Disinfestation and Inoculation with *Fusarium circinatum*—Determination of Inoculum Dose on/in the Seed

Pathogen free-seed of *P. radiata* from New Zealand was used both in seed contamination and artificial inoculation assays. Seeds were surface disinfested by immersion and shaking in: (1) 10% commercial bleach plus 0.1% Tween 20 for 15 min; (2) purified water (RIOs Type 3 RO, Millipore, Madrid, Spain) for 5 min, twice; (3) hydrogen peroxide 33% for 10 min; (4) sterile distilled water for 10 min, three times; and (5) dried in an air flow cabinet for 2 h.

Fusarium circinatum (CECT20759) was isolated from a *P. radiata* seedling in Guipuzkoa, Spain. This isolate was identified to be in vegetative compatibility group (VCG) A and Mating type 2 (Mat-2), the only mating type present in the Basque Country [25]. This isolate is stored on small pieces of sterile filter paper (Whatman no.1) at –20 °C. For inoculation purposes, it was cultured for 10 days at 22 °C on Potato Dextrose Agar (PDA) (Pronadisa, Madrid, Spain). The inoculum suspension was prepared by adding 10 mL of sterile purified water on the surface of a PDA plate where *F. circinatum* has grown for about 10 days at 22 °C, scraping with a spatula, and then filtering with glass wool washed

(Panreac Química, Barcelona, Spain). The spore concentration was determined using a haemocytometer and the suspension diluted to the desired concentration with sterile purified water. Inoculation was done by immersion and shaking at 150 rpm for 5 min in a spore suspension of *F. circinatum*. Seeds inoculated with water were used as control. Inoculated seeds were dried for 2 h in an air flow cabinet.

The viability of the inoculum was determined by plating dilution series onto PDA plates. The inoculum actually present in the seed after inoculation was estimated by grinding individually inoculated seeds (at various dose) in 1 mL sterile distilled water and plating on PDA plates. Also, when this estimate was below the detection limit, presence of inoculum was visualized growing the seeds on SFA (Selective Fusarium Agar) [26] plates.

2.3. Seed Contamination by *Fusarium circinatum* during Seed Storage

Differences during storage were tested at three conditions: 4 °C and 20% ± 3% RH, 20 °C and 25% ± 5% RH and 20 °C 65% ± 1% RH and at four sampling times. For each condition and sampling time, three plates with 101 surface disinfested *P. radiata* seeds per plate were used. One of these 101 seeds was inoculated by immersion in a suspension of 10³ spore/mL and plated on a 50 mm-diameter Petri dish with the other 100 non-inoculated seeds. Plates were incubated at the three conditions. Every week during four weeks, three plates of each condition were sampled and their seeds were plated on FSM to detect the number of seeds contaminated with *F. circinatum*.

2.4. In Vitro Effect of *Fusarium circinatum* on Seed Germination and Germling Growth

To assess the effects of inoculum location and doses on seed germination and germling growth, 50 surface-disinfested seeds of *P. radiata* were inoculated by immersion as described in Section 2.2, at doses of 10² or 10⁴ spores/mL: (1) protrusion-stage pregerminated seeds on water agar that mimic embryo, gametophyte and external infestation; (2) carborundum rubbed seeds to crack their coat and favor fungal penetration into seeds (imitating gametophyte and external infestation); and (3) no previous seed treatment to imitate external infestation of seed coats. Inoculation with water was done for each respective inoculation method. Each group of 50 seeds were germinated on a tray, covered with plastic bags to maintain high humidity and incubated at 22 °C with 12 h of light. Periodically until 48 days post inoculation (dpi), germlings were evaluated in a scale (0–2) for growth stage, where 0 = no germination, 1 = radicle present, and 2 = radicle and hypocotyl present; and for presence/absence of *F. circinatum* mycelia in stages 1 and 2.

Logit analysis was performed to test effects of inoculation method and inoculum dose on germination percentage at 20 dpi. Likewise, proportion of damaged germlings in growth stages 1 and 2 at the end of the experiment was analyzed by logistic regression on inoculation method and inoculum dose and their interaction.

2.5. Temporal Analysis of Pre- and Post- Emergence Damping-Off, and of Death of Established Seedlings

Pinus radiata surface-disinfested seeds were externally inoculated at doses of 10⁴ or 10⁶ spores/mL, stored at room temperature for one day until sowing in forest sowing 35-trays containing substrate composed of 80% peat plus 20% coconut fiber, and incubated at constant temperature 20 °C, and

12 h day/night light. Each treatment was applied to three trays, and trays were arranged in a randomized block design. The same conditions were repeated in a second experiment at the dose of 10^4 spores/mL.

There was a periodical monitoring and recording of seedling emergence and appearance of symptoms produced by *F. circinatum* seedling infection, such as girdling of root collar, wilting and die-back. Dead plants were sampled and some sections were plated on SFA medium to corroborate the presence of the pathogen. Identification of *F. circinatum* was confirmed morphologically on SNA [26] and PDA media based on morphological features in pure culture [27]. The last day of recording pre-emergence damping-off was at the date of the last emergence event. Post-emergence damping-off was recorded until 281 dpi, considering the first two months after emergence as damping-off and later on, damage of established seedlings. Analysis of variance was used to estimate the effect of inoculum dose on the proportion of pre- and post- emergence damping-off as well as proportion of dead plants at the end of the experiment. Data were transformed prior to the analysis using arc sine of the square root of the proportion. Mean values were compared by Student-Newman-Keuls multiple range tests.

Disease progress for the low inoculum dose (10^4 spores/mL) was regressed using different models (linear, monomolecular, logistic and Gompertz) [28]. The best model was chosen according to the lowest mean square error value (MSE) and the highest coefficient of determination value (R^2); then, disease infection rate was obtained from the slope parameter of the selected model [28]. Best fit of data was achieved using the linear model, that is, disease incidence regressed with time (dpi). Two lines were adjusted, one for post-emergence damping-off, and other for damage of established seedlings. The disease infection rate is the average of both replicates.

2.6. Persistence of *Fusarium circinatum* in Seedlings

As seedlings died from 77–281 dpi in the experiment described above, quantity of *F. circinatum* was estimated in a sample of them. Colony forming units (cfu) were calculated in five seedlings seed-inoculated with 10^6 and in 10 seedlings seed-inoculated with 10^4 spores/mL. Plants were surface disinfested by dipping them in 10% commercial bleach for 15 s, washed 3 times in sterile purified water, cut in pieces and ground in sterile purified water. Dilution series of these extracts were plated on SFA medium.

By the end of the same experiment (at 281 dpi), there were 30 seedlings apparently healthy. In order to further explore symptom development of these seedlings, they were transplanted to pots and grown in a greenhouse at 18 ± 5 °C with relative humidity of 55%–60%. Before this, seven of these seedlings were randomly chosen and processed as described above to estimate cfu of *F. circinatum*. The remaining 23 seedlings were monitored until 458 dpi. As seedlings died, plant parts (roots, stems and tips) were cut in transversal sections, plated on SFA and checked for presence of *F. circinatum*. Two plants survived at the end of this period.

3. Results

3.1. Natural Presence of *Fusarium circinatum* in/on Seeds of Symptomatic Trees

Total number of *P. radiata* seeds collected in eight plantations was 7100 from 71 trees. Out of these seeds, less than 1% (52 seeds from 12 trees) were infested with *F. circinatum* (mean of infested seeds = 0.73%, standard error \pm 0.244%). The pathogen was present on the coat of all infested seeds and

in some of them, *F. circinatum* was also infecting the gametophyte (mean percentage of infection 0.52 ± 0.219). In a small proportion, the embryo was also infected (0.10 ± 0.05). All seeds with an infected embryo had an infected gametophyte and seed coat as well (Figure 1).




Figure 1. Percentage of plantation-collected *Pinus radiata* (*P. radiata*) seeds with the presence of *F. circinatum* and their location on coat, gametophyte and embryo in relation to the tree disease severity (measured by the number of cankers) where seeds were collected.

Logistic regression analysis between percentage of infested seeds and tree disease severity (evaluated as number of cankers in a tree) showed a strong relationship (chi-square for wald test = 101.54, $df = 4$, p -value < 0.001). Odds ratio (OR) of number of infested seeds from trees with more than 1 canker compared to trees with only 1 canker was calculated from the logit model. According to this analysis, the odds that seeds would be infested was 40 times greater if seeds were collected from trees with five cankers (OR = 40.4, Wald chi-square < 0.001), and almost four times greater if seeds were collected from trees with two and three cankers (respectively, OR = 3.7, Wald chi-square = 0.0033; and OR = 3.9, Wald chi-square = 0.0064).

3.2. Seed Contamination by *Fusarium circinatum* during Storage

High relative humidity had a great effect on seed contamination at 20 °C, and this was true from the first week (Figure 2). There were no differences between number of contaminated seeds at 4 °C and 20 °C with low RH (20%). After four weeks, final percentage of contaminated seeds was 87% when incubated at 20 °C and 65% RH, while that value was 18.3% at 20 °C and 13.3% at 4 °C with low RH.




Figure 2. Percentage of *P. radiata* seeds contaminated by *F. circinatum* after incubating 100 disinfested seeds with one superficially inoculated seed. Incubation conditions were 20 °C with 25% or 65% of RH; and 4 °C and 20% RH. Each point is the average of three replicates. Bars are standard deviation.

3.3. In Vitro Effect of *Fusarium circinatum* on Seed Germination and Germling Growth

Germination of *P. radiata* seeds occurred until 20 dpi. At this date, percentage of total inoculated-seed germination was 86% (Figure 3). This value depended on inoculum dose (chi-square for wald test = 18.73, *df* = 2, *p*-value < 0.0001), and not on inoculation method (chi-square for wald test = 2.11, *df* = 2, *p*-value = 0.349), according to logit regression analysis of both factors. Germination mean was 78% ± 3.74% (standard error) for the dose of 10⁴ spores/mL, and 94% ± 1.41% for the dose of 10² spores/mL. This germination mean one did not differ significantly from the mean of untreated seeds (91% ± 0.82%) (Figure 3).




Figure 3. Total percentage germination of *P. radiata* seeds at 20 dpi for different inoculation doses and inoculation method: exp (4) = 10⁴ spores/mL; exp (2) = 10² spores/mL; E + IP = internal by immersion of pre-germinated seeds; E + IC = internal by immersion of seeds previously rubbed with carborundum; E = external by immersion. Treatments with different letters differ significantly at *P* = 0.05. No significant differences were among inoculation methods.

Total seed-inoculated germlings of *P. radiata* with only radicle grown (stage 1) at the end of the experiment (48 dpi) was 77 of total 300 seeds (equivalent to 26.3%) (Table 1). Of these, only two germlings did not show lesions or signs of infection by *F. circinatum* (Table 1). Final proportion of germlings in stage 1 depended on inoculation method (p -value = 0.0035) and on inoculum dose (p -value = 0.0001), but not on their interaction (p -value = 0.3).

According to the estimated odds ratio (OR) (Table 2), there were more germlings remaining in growth stage 1 when presence of *F. circinatum* was external than if the fungus was within the embryo (Figure 4) (in a proportion of 1 to 0.3; OR = 0.3, p -value = 0.001). There were not differences when the pathogen was external or infecting the gametophyte (p -value = 0.433). Regarding the dose, seeds inoculated with 10^4 spores/mL were three times more likely to remain in that stage than seeds inoculated with 10^2 spores/mL (Table 2). Proportion of damaged germlings at growth stage 2 depended on inoculation method (p -value = 0.0016) and dose (p -value < 0.0001). Estimated odds ratio (Table 2) showed that seeds with embryo and coat infected (protrusion-stage infection) were 2.4 times more likely to express symptoms than seeds with only coat infected (external infection).

Table 1. Number of *Pinus radiata* (*P. radiata*) germlings in each category of growth stage at 48 dpi, with presence or absence of *Fusarium circinatum* (*F. circinatum*) mycelia for each inoculation method and dose applied to surface-disinfested seeds.

Growth stage ⁽¹⁾	Fc ⁽²⁾	E + IP ⁽³⁾			E + IC ⁽³⁾			E ⁽³⁾			Sum	Sum of inoculated
		10 ⁴ ⁽⁴⁾	10 ²	wt	10 ⁴	10 ²	wt	10 ⁴	10 ²	wt		
2	0	0	4	43	1	3	40	0	2	46	139	10
2	1	32	38	0	10	37	0	19	33	0	169	169
Total 2		32	42	43	11	40	40	19	35	46	308	179
1	0	0	0	3	2	0	5	0	0	0	10	2
1	1	9	5	0	23	6	0	21	13	0	77	77
Total 1		9	5	3	25	6	5	21	13	0	87	79
Total 0		9	3	4	14	4	5	10	2	4	55	42
Total		50	50	50	50	50	50	50	50	50	450	300

⁽¹⁾ Growth stage: 0 = no germinated seed; 1 = germling with radicle; 2 = with hypocotyl and radicle; ⁽²⁾ Mycelia of *F. circinatum*: 1 = present; 0 = not present; ⁽³⁾ Inoculation method: E = external by immersion; IP = internal by immersion of pregerminated seeds; IC = internal by immersion of seeds previously rubbed with carborundum; ⁽⁴⁾ Inoculum dose: 10^4 spores/mL, 10^2 spores/mL, and wt = water as control.

Table 2. Odds ratio estimated by logistic regression of proportion of *P. radiata* damaged germlings on inoculation method and inoculum dose.

	Growth ¹ stage 1			Growth stage 2		
	OR	p -value	95% IC	OR	p -value	95% IC
IP ² vs. E	0.30	0.001	0.145, 0.612	2.35	0.0064	1.272, 4.351
IC vs. E	0.78	0.4328	0.421, 1.448	0.80	0.4536	0.443, 1.439
IP vs. IC	0.38	0.0095	0.185, 0.790	2.95	0.0006	1.589, 5.463
10^4 vs. 10^2	3.0	0.0001	1.707, 5.257	0.25	<0.0001	0.152, 0.411

¹ Growth stage of germling: 1= with radicle; 2 = with hypocotyl and radicle; ² Inoculation method: E = external by immersion; IP = internal by immersion of pregerminated seeds; IC = internal by immersion of seeds previously rubbed with carborundum.




Figure 4. Proportion of *P. radiata* germlings in each category of growth stage at 48 dpi, with presence of *F. circinatum* mycelia for each inoculation method and dose applied to surface-disinfested seeds. Exp (4) = 10⁴ spores/mL; exp (2) = 10² spores/mL; E + IP = internal by immersion of pre-germinated seeds; E + IC = internal by immersion of seeds previously rubbed with carborundum; E = external by immersion.

The percentage of germlings growing from inoculated seeds that developed radicle and hypocotyl at the end of the experiment (stage 2) was 60%. Of these germlings, all but 10 showed symptoms of *F. circinatum*, whereas germlings grown from non-inoculated seed showed no lesions (Table 1). The transmission rate from seed to germinating seeds and seedlings showing symptoms was determined and results are presented in Table 3. The rate of transmission from seed to germinating seeds causing pre-emergence death was always greater for the high inoculum dose whereas this transmission rate at post-emergence was always greater for the low dose. Infection after inoculation at the protrusion stage (E + IP) with 10⁴ spores/mL was lower than those rates for other methods at pre-emergence and higher at post-emergence.

Table 3. Transmission rate of *F. circinatum* from inoculated seeds to germinating seeds and symptomatic seedlings at 48 days after inoculation as determined *in vitro* and in potting soil assays.

Assay	Inoculation method ⁽¹⁾	Inoculum dose	Pre-emergence death	Post-emergence death	Total disease development	
"in vitro"	E + IP	10 ⁴	0.36 ⁽²⁾	0.64	1	
		10 ²	0.16	0.76	0.92	
	E + IC	10 ⁴	0.78	0.2	0.98	
		10 ²	0.2	0.74	0.94	
	E	10 ⁴	0.62	0.38	1	
		10 ²	0.3	0.66	0.96	
		average	10 ⁴	0.59	0.41	1
		average	10 ²	0.22	0.72	0.94
potting soil	E	10 ⁶	0.33	0.47	0.8	
		10 ⁴	0.15	0.13	0.28	

⁽¹⁾ Inoculation method: E = external by immersion; IP = internal by immersion of pregerminated seeds; IC = internal by immersion of seeds previously rubbed with carborundum; ⁽²⁾ For the *in vitro* assay, proportion of germlings with symptoms at growth stage 0 and 1.

3.4. Temporal Analysis of Pre- and Post- Emergence Damping-Off, and of Death of Established Seedlings

Inoculum on seeds was checked and it was present in all of them with estimates of 180–400 cfu/seed when inoculated with 10^6 spores/mL, and ≤ 10 cfu/seed when inoculum was 10^4 or 10^2 spores/mL. Highest emergence occurred respectively at 23 and 24 dpi for each repetition of the experiment, and disease symptoms became visible one week later. The major symptom was root collar rot, where mycelia could be observed. For older seedlings, the more generalized symptom was a wilting, starting at the top of the seedling and progressing down the stem until the seedling collapsed. Roots were underdeveloped.

Pre-emergence damping-off was 33.3% and 15.7% (Table 4) when seeds of *P. radiata* were inoculated respectively with 10^6 and 10^4 spores/mL, but only the higher dose had a significant effect compared to the low inoculum dose and the untreated control (Table 4). In post-emergence, inoculum dose had a significant effect on death of seedlings, with differences between both doses being greater in post-emergence (2-month-old seedlings) than in older seedlings (approximately 9-month-old seedlings) (Table 4).

Table 4. Effect of inoculum doses on percentage of pre- and post-emergence damping off, and death of established seedlings caused by *F. circinatum* on externally inoculated seeds of *P. radiata*.

Inoculum dose (spore/mL)	Pre-emergence damping-off (%)	Post-emergence damping-off (%) ⁽³⁾	Death of seedlings (%) ⁽³⁾
10^6 ⁽¹⁾	33.3 (4.67) ⁽²⁾ a ⁽⁴⁾	71.4 (5.07) a	88.4 (4.10) a
10^4 ⁽¹⁾	15.7 (3.10) b	24.7 (3.69) b	66.0 (5.73) b
0 (control)	7.6 (5.08) b	0	0

⁽¹⁾ Mean values of 3 replicates for the dose of 10^6 spore/mL, and of 6 replicates (three in each of 2 repetitions of the experiment) for the dose of 10^4 spore/mL; ⁽²⁾ In parenthesis, standard error of the mean value; ⁽³⁾ Post-emergence damping-off was recorded for seedlings at 2 months after emergence, and for established plants at 281 dpi; ⁽⁴⁾ Numbers followed by the same letter within a column are not significantly different (significant value 0.05) according to Student-Newman-Keuls multiple range tests.

Disease incidence progressed throughout the experiment (until 281 dpi) with two peaks for which disease rate was highest (Figure 5). One peak was during post-emergence stage from 30–36 dpi, and the other was during the period of established seedlings, at 93 dpi for the first repetition, and at 114 dpi for the second one. (Figure 5). Although disease rate was not uniform in each of the two periods, the best model describing the disease progress curve was the lineal model (disease incidence regressed on time) being the slope of that line an estimate of the disease infection rate. The rate of disease incidence per day was 0.0025 ± 0.00067 (standard error) for damping-off in post-emergence and 0.0016 ± 0.00024 during the period of established seedlings. However, these rates were not significantly different.




Figure 5. Disease progress curve on seedlings of *P. radiata* grown from seeds externally inoculated with 10^4 spores/mL of *F. circinatum*. Each curve is the mean of three replicates in each experiment repetition (repetition 1 for orange curve and repetition 2 for blue curve). Dpi means days post inoculation.

Specific values of disease incidence at 48 dpi were used to determine the transmission rate from seed to symptomatic seedlings (Table 3). Transmission rates at this time of the disease development are similar in pre- and post-emergence for the low inoculum dose (10^4 spores/mL), and higher in post-emergence for the high inoculum dose.

3.5. Persistence of *Fusarium circinatum* in Seedlings

Fusarium circinatum was quantified in a sample of *P. radiata* dead seedlings (five and 10 seedlings from seeds inoculated respectively with 10^6 and 10^4 spores/mL). Populations in roots were in the range of 10^5 – 10^7 cfu/gr fresh weight for both seed inoculation doses, and in stems between 10^2 and 10^5 cfu/gr fresh weight. *F. circinatum* was not detected in the stem of three dead plants, however, it could be present since detection level was about 120 cfu/gr. Estimates of *F. circinatum* cfu were in those ranges regardless of whether seedlings came from inoculated seeds with either 10^6 or 10^4 as it was observed in root populations.

At 281 dpi, 30 seedlings remained symptomless. On seven of these, *F. circinatum* was estimated and it was detected in roots of five plants, with populations up to 10^4 cfu/gr. In stems of these plants, *F. circinatum* was only detected in one plant (10^5 cfu/gr). The other 23 seedlings were moved to a greenhouse and grown in pots from 281 dpi to 479 dpi. All plants (from both seed inoculation doses) except two of them died in this period, and *F. circinatum* was detected in all dead plants, in roots and stems, except in the stem of one of these plants. The two plants remaining at the end of the experiment were symptomless; however, *F. circinatum* was detected in their roots, stems and tips. The final percentage of symptomless seedlings at 281 dpi was 14% and at 481 dpi was 1%.

4. Discussion

F. circinatum was isolated from *P. radiata* seeds from trees with cankers of pitch canker disease, but at a frequency less than 1%. This is a very low rate compared to previous studies. In California,

F. circinatum was isolated from up to 83% of seeds collected from *P. radiata* cones on infected branches [11]. For other *Pinus* species, the pathogen was also isolated from 98% of *P. palustris* seeds [29] of which 84% was internal infection; in *P. taeda* seeds, external presence varied from 7%–61%, and internal infection from 1%–34% [30]; in *P. elliottii* seeds, internal presence of *F. circinatum* was up to 30% [15].

Besides *Pinus* species, environmental conditions also influence the extent of infection [31]. Seeds analyzed here were collected in one year from eight different locations, and this short sampling time could influence the low level of *F. circinatum* detected. However, these results are in agreement with those obtained in routine seed health testing that has been performed in Spain since *F. circinatum* was detected in 2005, which are very low (personal communication). In this case, it is assumed that seeds are collected from orchard trees with no pitch canker symptoms, but *F. circinatum* may also be present on seeds collected from cones of apparently healthy branches [11] coming from airborne spores that enter the cone when it is open.

The low seed infection rate could be also due to the location of sampled cones, which may be on branches without symptoms, especially in those trees sampled with one or two cankers. For these trees, percentage of seeds with presence of *F. circinatum* was less than 1% for each plantation (Figure 1), but again this is very low compared to reports from two locations in California whereby 9.2% and 11.7% of seeds were infected from symptomless branches [11]. Another reason to explain the low infection could be the method used here for detecting the pathogen on seeds. Plating seeds prior to germination does not detect cryptic infections, which are only detected from emerging seedlings [11].

Fusarium circinatum was mainly on the seed coat and less frequently in the gametophyte. This reveals that infection, at least mainly, is not systemically transmitted from the mother plant to the embryo, but is transmitted indirectly by air-borne spores entering cones at time of pollination or/and penetrating mature cones through small cracks. No coincidence of pollination and dispersal of fungal spores or low amount of them could explain the low seed infection found.

The present study showed that external presence of the pathogen on the seed can increase due to contamination during storage, and then seeds may become an important inoculum source in nurseries. Management of storage conditions, especially low relative humidity level, can help to prevent contamination and therefore it is highly recommended to store seeds under low relative humidity. Further studies are required to optimize storage conditions, and to determine the best treatments to reduce or eliminate the external presence of the fungus, or at least to reduce chances of contamination.

Our work showed that transmission from seed to seedling is accomplished even with presence of *F. circinatum* limited to the seed coat (external inoculation) with an inoculum dose lower than 10 spores per seed, and these seeds can be a source of primary inoculum for pitch canker disease in nurseries. The fungus was transmitted to seedlings causing death in pre- or post- emergence at different rates depending on the inoculum dose. Rates of transmission were determined from seeds to symptomatic seedlings, but we showed that fungus was also transmitted to the remaining symptomless seedlings in potting soil. Of these, some developed disease later and others did not, but the fungus was detected in all seedlings coming from seeds originally inoculated with *F. circinatum*. Seed transmission rate was higher *in vitro* than in potting soil, since the humid environment created in germination trays was more conducive for fungus growth and disease development. This suggests that humidity level in soil during seed germination may influence pathogen transmission in nurseries.

When the disease outbreak occurs in nurseries, the pathogen could be disseminated to new plantations through symptomless seedlings. Seedlings progressively developed pitch canker symptoms during the first 15 months, but a few did not develop symptoms at the end of the experiment (two of a total 203 seedlings that reached post-emergence). It is unknown if these established seedlings will show disease once grown in a plantation, but this could be one of the reasons why pitch canker disease remains latent for years, as has been observed in South Africa [32] and in Spain. The pathogen was detected in the roots of both types of seedlings, with and without symptoms, as was previously reported by Storer *et al.* [11]. We did not detect the pathogen in two symptomless seedlings (inoculated with low dose) but this could be due to either *F. circinatum* population being less than the detection level (13–20 cfu/mL) or simply an infection escape. We found that fungus population present in the roots of seedlings with symptoms was always higher than in seedlings without symptoms, this population being lower than 10^4 cfu/g. Pathogen population in stems of the seedlings with symptoms was always lower than in roots. Because studies on colonization and symptoms of *F. circinatum* in established plants have focused on the aerial part of the tree (see e.g., [33,34]), the role of root colonization on persistence and dissemination of *F. circinatum* may have been underestimated.

Pre-emergence damping-off is caused by germination failure or damages to the radicle. Results from the germling growth assay indicated that final germination *in vitro* was not affected by whether the inoculum was located in embryo, gametophyte or coat, as suggested when comparing different inoculation methods (Figure 3). On the contrary, damages in the radicle preventing further growth were caused by presence of *F. circinatum* in the gametophyte or on the seed coat (Table 3). The reason why there was less damage when the embryo was infected is unknown, but could be influenced by the time the embryo was infected, in the protrusion stage, which may not stop seeds from growing. It may suggest that *F. circinatum* inhibited hypocotyl development since the number of germlings that remained in stage 1 after inoculation at the high dose was about double in the treatment without protrusion pretreatment. Higher inoculum dose led to greater damage at the germination plus radicle stage (OR = 3, Table 2) than at the stage of radicle plus hypocotyl (OR = 0.25, Table 2). This is in agreement with damages quantified in pre-emergence, which was only significant for seeds inoculated with a high dose (Table 3).

Once seedlings emerged, high inoculum doses caused greater mortality in the first two months, while low dose caused damage that progressed steadily until 281 dpi (Table 3, Figure 5). Viljoen *et al.* [3] observed that mortality of three-year-old seedlings of *P. patula* was lower than that of newly germinated seedlings. So, our results corroborate a lower mortality for established seedlings, but only for high inoculum doses. We determined the infection rate in optimal conditions for the pitch canker disease at low inoculum doses, during post-emergence and established seedlings, from lineal models adjusted to the disease progress curve during each growth period (Figure 5). Disease infection rates were not significantly different, corroborating that mortality in post-emergence and in established seedlings is similar when seeds were inoculated with 10^4 spores/mL.

5. Conclusions

In Spain, *P. radiata* is an exotic species growing only in plantations with seedlings coming from nurseries. We showed that transmission of *F. circinatum* from symptomatic trees to seeds is extremely low compared to other pine species and locations. The pathogen is located mainly on the seed coat, and

could contaminate other seeds during storage. This inoculum can be transmitted to new seedlings causing death at pre- and post-emergence. Very low presence of *F. circinatum* as low as 10 spores or less per seed (inoculum on the coat when seeds inoculated with 10^4 spores/mL) may transmit disease with a rate of up to 90% to established seedlings causing their death. However, a small proportion of seedlings may stay symptomless but latent-infected. Approximately one year old *P. radiata* seedlings are moved from nurseries to new plantations. We have determined that the proportion of symptomless seedlings with latent infection is 3.6% when seedlings are 365 days old and 1% when they are 481 days old. Dissemination disease risk is higher for low pathogen inoculum since it will take more time for seedlings to show symptoms although they are latent-infected, and, therefore, a higher proportion of these seedlings will be taken to the new plantations.

Acknowledgments

This study was supported by Ministerio de Economía y Competitividad (Subprograma de Proyectos de I + D orientado a los Recursos y Tecnologías Agrarias en coordinación con las Comunidades Autónomas) Project RTA 2012-00015.

Author Contributions

Eugenia Iturriza developed conceptual ideas, designed the study, conducted measurements and data analysis in field experiments. Margarita Elvira-Recuenco and Rosa Raposo developed conceptual ideas, design the study, conducted measurements and data analysis in lab experiments, and wrote the paper.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Hepting, G.H.; Roth, E.R. Pitch canker, a new disease of some southern pines. *J. For.* **1946**, *44*, 742–744.
2. Wingfield, M.J.; Hammerbacher, A.; Ganley, R.J.; Steenkamp, E.T.; Gordon, T.R.; Wingfield, B.D.; Coutinho, T.A. Pitch canker caused by *Fusarium circinatum*—A growing threat to pine plantations and forests worldwide. *Australas. Plant Pathol.* **2008**, *37*, 319–334.
3. Viljoen, A.; Wingfield, M.J.; Marasas, W.F.O. First report of *Fusarium subglutinans* f. sp. *pini* on seedlings in South Africa. *Plant Dis.* **1994**, *78*, 309–312.
4. Wingfield, M.J.; Jacobs, A.; Coutinho, T.A.; Ahumada, R.; Wingfield, B.D. First report of the pitch canker fungus, *Fusarium circinatum*, on pines in Chile. *Plant Pathol.* **2002**, *51*, 397–397.
5. Landeras, E.; Garcia, P.; Fernandez, Y.; Brana, M.; Fernandez-Alonso, O.; Mendez-Lodos, S.; Perez-Sierra, A.; Leon, M.; Abad-Campos, P.; Berbegal, M.; *et al.* Outbreak of pitch canker caused by *Fusarium circinatum* on *Pinus* spp. in Northern Spain. *Plant Dis.* **2005**, *89*, 1015–1015.
6. Braganca, H.; Diogo, E.; Moniz, F.; Amaro, P. First Report of Pitch Canker on Pines Caused by *Fusarium circinatum* in Portugal. *Plant Dis.* **2009**, *93*, 1079.

7. Pfenning, L.H.; Costa, S.D.S.; de Melo, M.P.; Costa, H.; Ventura, J.A.; Auer, C.G.; dos Santos, A.F. First report and characterization of *Fusarium circinatum*, the causal agent of pitch canker in Brazil. *Trop. Plant Pathol.* **2014**, *39*, 210–216.
8. Carey, W.A.; Oak, S.W.; Enebak, S.A. Pitch canker ratings of longleaf pine clones correlate with *Fusarium circinatum* infestation of seeds and seedling mortality in containers. *For. Pathol.* **2005**, *35*, 205–212.
9. Coutinho, T.A.; Steenkamp, E.T.; Mongwaketsi, K.; Wilmot, M.; Wingfield, M.J. First outbreak of pitch canker in a South African pine plantation. *Australas. Plant Pathol.* **2007**, *36*, 256–261.
10. Berbegal, M.; Perez-Sierra, A.; Armengol, J.; Gruenwald, N.J. Evidence for Multiple Introductions and Clonality in Spanish Populations of *Fusarium circinatum*. *Phytopathology* **2013**, *103*, 851–861.
11. Storer, A.J.; Gordon, T.R.; Clark, S.L. Association of the pitch canker fungus, *Fusarium subglutinans* f. sp. *pini*, with Monterey pine seeds and seedlings in California. *Plant Pathol.* **1998**, *47*, 649–656.
12. Huang, J.W.; Kuhlman, E.G. Fungi associated with damping-off of slash pine-seedlings in Georgia. *Plant Dis.* **1990**, *74*, 27–30.
13. Barrowsbroaddus, J.; Dwinell, L.D. Branch dieback and cone and seed infection caused by *Fusarium moniliforme* var *subglutinans* in a loblolly pine seed orchard in South Carolina. *Phytopathology* **1985**, *75*, 1104–1108.
14. Anderson, R.L. New method for assessing contamination of slash and loblolly-pine seeds by *Fusarium-moniliforme* var *subglutinans*. *Plant Dis.* **1986**, *70*, 452–453.
15. Anderson, R.L.; Belcher, E.; Miller, T. Occurrence of seed fungi inside slash pine seeds produced in seed orchards in the United-States. *Seed Sci. Technol.* **1984**, *12*, 795–799.
16. Bloomberg, W.J. The epidemiology of forest nursery diseases. *Annu. Rev. Phytopathol.* **1985**, *23*, 83–96.
17. Perez-Sierra, A.; Landeras, E.; Leon, M.; Berbegal, M.; Garcia-Jimenez, J.; Armengol, J. Characterization of *Fusarium circinatum* from *Pinus* spp. in northern Spain. *Mycol. Res.* **2007**, *111*, 832–839.
18. Iturrutxa, E.; Mesanza, N.; Elvira-Recuenco, M.; Serrano, Y.; Quintana, E.; Raposo, R. Evaluation of genetic resistance in *Pinus* to pitch canker in Spain. *Australas. Plant Pathol.* **2012**, *41*, 601–607.
19. Iturrutxa, E.; Ganley, R.J.; Raposo, R.; Garcia-Serna, I.; Mesanza, N.; Kirkpatrick, S.C.; Gordon, T.R. Resistance levels of Spanish conifers against *Fusarium circinatum* and *Diplodia pinea*. *For. Pathol.* **2013**, *43*, 488–495.
20. Ios, R. Detection of *Gibberella Circinata* on *Pinus* spp. (Pine) and *Pseudotsuga Menziesii* (Douglas-Fir) Seed. In *International Rules for Testing*, Proceedings of Ordinary General Meeting, Edinburgh, United Kingdom, 2014; ISTA: Bassersdorf, Switzerland, 2015.
21. Inman, A.R.; Kirkpatrick, S.C.; Gordon, T.R.; Shaw, A.V. Limiting effects of low temperature on growth and spore germination in *Gibberella circinata*, the cause of pitch canker in pine species. *Plant Dis.* **2008**, *92*, 542–545.
22. Gordon, T.R.; Okamoto, D.; Storer, A.J.; Wood, D.L. Susceptibility of five landscape pines to pitch canker disease, caused by *Fusarium subglutinans* f. sp. *pini*. *HortScience* **1998**, *33*, 868–871.
23. Aitken, J.; Iturrutxa, E. *Disease-Free Import of Pinus Seed*; Report to MAF-For; The Tree Lab: Biosecurity, New Zealand, 30 September 2004.

24. Aegerter, B.J.; Gordon, T.R. Rates of pitch canker induced seedling mortality among *Pinus radiata* families varying in levels of genetic resistance to *Gibberella circinata* (anamorph *Fusarium circinatum*). *For. Ecol. Manag.* **2006**, *235*, 14–17.
25. Iturrutxa, E.; Ganley, R.J.; Wright, J.; Heppe, E.; Steenkamp, E.T.; Gordon, T.R.; Wingfield, M.J. A genetically homogenous population of *Fusarium circinatum* causes pitch canker of *Pinus radiata* in the Basque Country, Spain. *Fungal Biol.* **2011**, *115*, 288–295.
26. Leslie, J.F.; Summerell, B.A. *The Fusarium Laboratory Manual*, 3rd ed.; Blackwell: Ames, IA, USA, 2006; p. 370.
27. EPPO. *Gibberella circinata*. *Bull. OEPP/EPPO Bull.* **2009**, *39*, 298–309.
28. Campbell, C.L.; Madden, L.V. *Introduction to Plant Disease Epidemiology*; Wiley: Hoboken, NJ, USA, 1990; p. 532.
29. Runion, G.B.; Bruck, R.I. Effects of Thiabendazole-DMSO treatment of longleaf pine seed contaminated with *Fusarium subglutinans* on germination and seedling survival. *Plant Dis.* **1988**, *72*, 872–874.
30. Barrowsbroaddus, J. Colonization of cones and seed of loblolly-pine following inoculation with *Fusarium subglutinans*. *Plant Dis.* **1990**, *74*, 1002–1005.
31. Dwinell, L.D.; Barrowsbroaddus, J.B.; Kuhlman, E.G. Pitch canker—A disease complex of Southern pines. *Plant Dis.* **1985**, *69*, 270–276.
32. Van Wyk, S.J.P.; Boutigny, A.-L.; Coutinho, T.A.; Viljoen, A. Sanitation of a South African Forestry Nursery Contaminated with *Fusarium circinatum* Using Hydrogen Peroxide at Specific Oxidation Reduction Potentials. *Plant Dis.* **2012**, *96*, 875–880.
33. Correll, J.C.; Gordon, T.R.; McCain, A.H.; Fox, J.W.; Wood, D.L.; Schultz, M.E. Pitch canker in California: pathogenicity, distribution, and canker development on Monterey pine (*Pinus radiata*). *Plant Dis.* **1991**, *75*, 676–682.
34. Storer, A.J.; Wood, D.L.; Gordon, T.R. The epidemiology of pitch canker of Monterey pine in California. *For. Sci.* **2002**, *48*, 694–700.